

# General Properties Of Dalitz Plots

GlueX-doc-181-v0

Curtis A. Meyer  
Carnegie Mellon University

July 2004

## Abstract

This document derives some general properties of Dalitz plots. In particular, a derivation of the relation between the density along a decay band and the cosine of the decay angles is presented. It is also shown that given two identical particles, how the helicity angles of the two decay chains are related.

## 1 Properties of Dalitz Plots

We will consider the Dalitz plot as evaluated in the center of mass frame of the three daughter particles. Under that assumption, we have the following constraints:

$$M = E_1 + E_2 + E_3 \quad (1)$$

$$0 = \vec{p}_1 + \vec{p}_2 + \vec{p}_3 \quad (2)$$

$$E_1 = \sqrt{p_1^2 + m_1^2} \quad (3)$$

$$E_2 = \sqrt{p_2^2 + m_2^2} \quad (4)$$

$$E_3 = \sqrt{p_3^2 + m_3^2}. \quad (5)$$

The Dalitz plot is made by plotting either the kinetic energy of one particle versus another, *e.g.*  $T_3$  versus  $T_2$ , or by plotting the invariant mass of one pair of particles versus another pair, *e.g.*  $m_{13}^2$  versus  $m_{12}^2$ .

The invariant mass of a pair of particles is written as:

$$m_{ij}^2 = (E_i + E_j)^2 - (\vec{p}_i + \vec{p}_j)^2 \quad (6)$$

$$= m_i^2 + m_j^2 + 2(E_i E_j - \vec{p}_i \cdot \vec{p}_j) \quad (7)$$

$$= m_i^2 + m_j^2 + 2(E_i E_j - p_i p_j \cos \theta_{ij}). \quad (8)$$

There is also a global constraint relating the seven masses in the problem:

$$M^2 = m_{12}^2 + m_{13}^2 + m_{23}^2 - m_1^2 - m_2^2 - m_3^2. \quad (9)$$

Consider now the addition of  $m_{12}^2$  and  $m_{13}^2$ , where we replace  $\vec{p}_3 = -\vec{p}_1 - \vec{p}_2$  and  $E_3 = M - E_1 - E_2$ . It is easily shown that

$$E_1 = \frac{M^2 + m_1^2 - m_{23}^2}{2M} \quad (10)$$

$$E_2 = \frac{M^2 + m_2^2 - m_{13}^2}{2M} \quad (11)$$

$$E_3 = \frac{M^2 + m_3^2 - m_{12}^2}{2M}. \quad (12)$$

This confirms that the original statement on energy and invariant-mass squared being equivalent.

Now let us consider a situation where our particle  $M$  decays into  $m_{23}$  recoiling against particle 1. We want to define a coordinate system whose  $z$ -axis is along  $\vec{p}_1$  and whose  $y$ -axis is normal to the decay plane of the three particles. We then want to boost along  $\hat{z}$  into the rest frame of  $m_{23}$  and look at the angle between particle 2 and the  $\hat{z}$  axis in that frame,  $\cos \theta_2$ . In the rest frame of  $M$ , the momentum of particle 2 is given as

$$\vec{p}_2 = \begin{pmatrix} p_2 \sin \theta_{12} \\ 0 \\ p_2 \cos \theta_{12} \end{pmatrix} \quad (13)$$

where  $\theta_{12}$  is the smallest angle between  $\vec{p}_1$  and  $\vec{p}_2$ . The Lorentz boost into the rest frame of  $m_{23}$  (called the helicity frame) is given by :

$$\beta = -\frac{p_1}{E_1} \quad (14)$$

$$\gamma = \frac{E_1}{m_1} \quad (15)$$

$$\beta\gamma = -\frac{p_1}{m_1} \quad (16)$$

This then yields that

$$E_2^h = \gamma E_2 - \beta\gamma p_2 \cos \theta_{12} \quad (17)$$

$$\vec{p}_2^h = \begin{pmatrix} p_2 \sin \theta_{12} \\ 0 \\ \gamma p_2 \cos \theta_{12} - \beta\gamma E_2 \end{pmatrix} \quad (18)$$

For a given  $m_{23}$ , the magnitude of  $\vec{p}_2^h$  and  $E_2^h$  are given as:

$$p_2^h = \sqrt{\frac{(m_{23}^2 + m_3^2 - m_2^2)^2}{4m_{23}^2} - m_3^2} \quad (19)$$

$$E_2^h = \sqrt{\frac{(m_{23}^2 - m_3^2 + m_2^2)^2}{4m_{23}^2}} \quad (20)$$

Now, we can write down what the  $\cos \theta_2$  is in the helicity frame. This can be simplified by inserting  $\beta, \gamma$  from above and noting that  $p_2 \cos \theta_{12}$  can be replaced by:  $(m_1^2 + m_2^2 + 2E_1 E_2 - m_{12}^2)/p_1$

$$\cos \theta_2 = \frac{\gamma p_2 \cos \theta_{12} - \beta\gamma E_2}{p_2^h} \quad (21)$$

$$= \frac{1}{p_2^h} \left[ \frac{E_1}{m_1} \frac{m_1^2 + m_2^2 + 2E_1 E_2 - m_{12}^2}{p_1} - \frac{p_1}{m_1} E_2 \right] \quad (22)$$

$$= \frac{1}{p_2^h p_1 m_1} [E_1(m_1^2 + m_2^2 + 2E_1 E_2 - m_{12}^2) - p_1^2 E_2] \quad (23)$$

$$= \frac{1}{p_2^h p_1 m_1} [E_1(m_1^2 + m_2^2 - m_{12}^2) + m_1^2 E_2] \quad (24)$$

$$= \frac{1}{p_2^h p_1 m_1} \left[ E_1(m_1^2 + m_2^2 - m_{12}^2) + m_1^2 \frac{M^2 + m_2^2 - m_{13}^2}{2M} \right] \quad (25)$$

$$= \frac{[(M^2 + m_1^2 - m_{23}^2)(m_1^2 + m_2^2 - m_{12}^2) + m_1^2(M^2 + m_2^2 - m_{13}^2)]}{p_2^h p_1 m_1 2M} \quad (26)$$

$$= \frac{(M^2 + m_1^2 - m_{23}^2)(m_{23}^2 + m_{13}^2 - M^2 - m_3^2) + m_1^2(M^2 + m_2^2 - m_{13}^2)}{p_2^h p_1 m_1 2M} \quad (27)$$

$$= \frac{m_{13}^2 [M^2 - m_{23}^2] + [(m_1^2 - (m_{23}^2 - M^2))((m_{23}^2 - M^2) - m_3^2) + m_1^2(M^2 + m_2^2)]}{p_2^h p_1 m_1 2M} \quad (28)$$

For a fixed value of  $m_{23}^2$ ,  $p_1$ , and  $p_2^h$  are also constants. This means that the value of  $\cos \theta_2$  is a linear function of  $m_{13}^2$ .

$$\cos \theta_2 = \frac{m_{13}^2 [M^2 - m_{23}^2] + [(m_1^2 - (m_{23}^2 - M^2))((m_{23}^2 - M^2) - m_3^2) + m_1^2(M^2 + m_2^2)]}{(m_1/2m_{23})\sqrt{[(M^2 + m_1^2 - m_{23}^2)^2 - 4M^2m_1^2][(m_{23}^2 + m_3^2 - m_2^2)^2 - 4m_{23}^2m_3^2]}} \quad (29)$$

## 2 Symmetries

Let us now assume that  $m_2 = m_3 = m$ . This means that we should expect symmetries around the line  $m_{12}^2 = m_{13}^2$  in the plot. Let us explore this to see what we find. We will define two possible decay chains here:

$$X \rightarrow A_{12}3 \rightarrow 123 \quad (30)$$

$$X \rightarrow B_{13}2 \rightarrow 132. \quad (31)$$

In decay chain 30, we have four angles:  $(\theta_{gj}^A, \phi_{gj}^A, \theta_h^A, \phi_h^A)$ , while in decay chain 31, we have four angles:  $(\theta_{gj}^B, \phi_{gj}^B, \theta_h^B, \phi_h^B)$ . Using equation 29 and assuming first that  $\vec{p}_3$  defines the  $\hat{z}$  axis, we find that

$$\cos \theta_h^A = \frac{m_{23}^2 [M^2 - m_{12}^2] + [(m^2 - (m_{12}^2 - M^2))((m_{12}^2 - M^2) - m^2) + m^2(M^2 + m_1^2)]}{(m/2m_{12})\sqrt{[(M^2 + m^2 - m_{12}^2)^2 - 4M^2m^2][(m_{12}^2 + m^2 - m_1^2)^2 - 4m_{12}^2m^2]}} \quad (32)$$

Now, if we choose  $\vec{p}_2$  to define the  $\hat{z}$  axis, then we find that the angle of particle 1 in the helicity frame is:

$$\cos \theta_h^B = \frac{m_{23}^2 [M^2 - m_{13}^2] + [(m^2 - (m_{13}^2 - M^2))((m_{13}^2 - M^2) - m^2) + m^2(M^2 + m_1^2)]}{(m/2m_{13})\sqrt{[(M^2 + m^2 - m_{13}^2)^2 - 4M^2m^2][(m_{13}^2 + m^2 - m_1^2)^2 - 4m_{13}^2m^2]}}. \quad (33)$$

Finally, if we look along the symmetry axis where  $m_{12}^2 = m_{13}^2$ , then these two helicity angles are exactly equal. Now let us assume that in the rest frame of  $X$ , we have some coordinate system whose unit vectors are given as  $(\hat{e}_x, \hat{e}_y, \hat{e}_z)$ . If we now use these directions to define a  $y$ -axis in the helicity frame as  $\hat{y}$  is along  $\hat{e}_z \times \hat{z}$ . The question is now what is  $\phi_h$  in the two frames? This angle is actually just the angle between the normal to the decay plane and the  $\hat{y}$  axis as defined above. The normal to the decay plane is given as follows, where the angle is measured in the rest frame of  $X$ .

$$\hat{n} = \frac{\vec{p}_1 \times \vec{p}_2}{p_1 p_2 \sin \theta_{12}} \quad (34)$$

$$= \frac{\vec{p}_2 \times \vec{p}_3}{p_2 p_3 \sin \theta_{23}} \quad (35)$$

$$= \frac{\vec{p}_3 \times \vec{p}_1}{p_3 p_1 \sin \theta_{31}} \quad (36)$$

The  $y$ -axis is given as:

$$\hat{y}^A = \frac{\vec{p}_3 \times \hat{e}_z}{p_3 \sin \theta_{3z}} \quad (37)$$

$$\hat{y}^B = \frac{\vec{p}_2 \times \hat{e}_z}{p_2 \sin \theta_{2z}} \quad (38)$$

where again, the angle is measured in the rest-frame of  $X$ . The angle  $\phi$  can now be determined as  $\cos \phi = \hat{n} \cdot \hat{y}$ .

$$\cos \phi_h^A = \left( \frac{\vec{p}_2 \times \vec{p}_3}{p_2 p_3 \sin \theta_{23}} \right) \cdot \left( \frac{\vec{p}_3 \times \hat{e}_z}{p_3 \sin \theta_{3z}} \right) \quad (39)$$

$$= \frac{(\vec{p}_2 \cdot \vec{p}_3)(\vec{p}_3 \cdot \hat{e}_z) - (\vec{p}_2 \cdot \hat{e}_z)(\vec{p}_3 \cdot \vec{p}_3)}{p_2 p_3^2 \sin \theta_{23} \sin \theta_{3z}} \quad (40)$$

Similarly, we can write that:

$$\cos \phi_h^B = \frac{(\vec{p}_2 \cdot \vec{p}_2)(\vec{p}_3 \cdot \hat{e}_z) - (\vec{p}_2 \cdot \hat{e}_z)(\vec{p}_3 \cdot \vec{p}_2)}{p_2^2 p_3 \sin \theta_{23} \sin \theta_{2z}}. \quad (41)$$

It should be noted that both of these formulas have terrible pathologies. If any of the sin functions are zero, or if either  $p_2$  or  $p_3$  are zero. Other than this, there should be an exact one-to-one relationship between  $\phi_h^A$  and  $\phi_h^B$ . Let's now restrict ourselves to the diagonal. Here we have that  $m_2 = m_3 = m$ ,  $p_2 = p_3 = p$ ,  $E_2 = E_3 = E$  and  $\cos \theta_{12} = \cos \theta_{13}$ . This means that  $\theta_{12} = \theta_{13} = \theta$ .

$$\cos \phi_h^A = \frac{\cos \theta_{23} \cos \theta_{3z} - \cos \theta_{2z}}{\sin \theta_{23} \sin \theta_{3z}} \quad (42)$$

$$\cos \phi_h^B = \frac{\cos \theta_{3z} - \cos \theta_{23} \cos \theta_{2z}}{\sin \theta_{23} \sin \theta_{2z}} \quad (43)$$

Finally, we should note that a given point in the Dalitz plot specifies the magnitudes of the three 3-momenta in the decay plane. It does **not** specify the orientation of the decay plane. We can apply any arbitrary rotation to the three particles we want, and we will still get the same entry in the Dalitz plot. This means that a particular point in the Dalitz plot does not translate into a unique  $\theta_{gj}$  or  $\phi_{gz}$  as it does for the angles in the helicity frame.