

Hall- D and the GlueX Experiment at Jefferson Lab

Simon Taylor / JLAB

Exotic Mesons

The 12 GeV Upgrade

Hall D

GlueX

Outlook

Gluonic Degrees of Freedom

Perturbative

Non-Perturbative

Asymptotic Freedom



Confinement

Small Distance
High Energy

Large Distance
Low Energy

High Energy
Scattering

Gluon
Jets
Observed



Gluonic Degrees of Freedom

Perturbative



Asymptotic Freedom

High Energy Scattering

Gluon Jets Observed



Small Distance
High Energy




Non-Perturbative

Confinement


Large Distance
Low Energy

Spectroscopy

lepton
meson
baryon



Gluonic
Degrees of Freedom
Missing



Quark Pairs and Triplets and Glue?

- Conventional hadrons: $q\bar{q}$ or qqq
 - Strong force mediated by gluons...
... but glue not needed to describe these states (quark model)...
- Gluons carry color charge → can couple to each other

Allowed systems: gg , ggg , $q\bar{q}g$, $q\bar{q}q\bar{q}$


 $\underbrace{gg}_{\text{Glueballs}}, \underbrace{ggg}_{\text{Hybrids}}, \underbrace{q\bar{q}g, q\bar{q}q\bar{q}}_{\text{Molecules}}$
Gluonic Excitations

Excitation of glue can lead to exotic quantum numbers J^P not possible in simple quark model

Quark Pairs and Triplets and Glue?

- Conventional hadrons: $q\bar{q}$ or qqq
 - Strong force mediated by gluons...
... but glue not needed to describe these states (quark model)...
- Gluons carry color charge → can couple to each other


Allowed systems: gg , ggg , $q\bar{q}g$, $q\bar{q}q\bar{q}$

$\underbrace{\hspace{10em}}$ $\underbrace{\hspace{10em}}$ $\underbrace{\hspace{10em}}$

Glueballs Hybrids Molecules



Gluonic Excitations

*Excitation of glue can lead to **exotic quantum numbers** J^P not possible in simple quark model*




GlueX Focus: light- quark mesons

Lattice Calculations



D. Leinweber

Flux tubes realized in LQCD




G. Bali


Plucking the Flux Tube

How do we look for gluonic degrees of freedom in spectroscopy?

Normal meson:
flux tube in
ground state
 $m=0$
 $CP=(-1)^S$




Hybrid meson:
flux tube in
excited state
 $m=1$
 $CP=(-1)^S$




Nonets characterized by given J^{PC}


$B^+ \bar{D}^- (\ell^-) SS$




In the first-excited state we have two degenerate transverse modes with $J=1^-$ - clockwise and counter-clockwise - and their linear combinations lead to $J^{PC}=1^{-+}$ or $J^{PC}=1^{+-}$ for the excited flux-tube




Meson Map




Meson Map



Meson Map






Meson Map




Production of Exotic Mesons

- Choice of probe may determine accessible quantum numbers
 - Photon beam increases chance for producing mesons with exotic quantum numbers



QCD Exotic Topologies

The GlueX Detector Design has been driven by the need to carry out Amplitude analysis.



$$\begin{array}{ccc} \pi_1 \eta_1 \eta_1 & b_2 h_2 h_2 & b_0 h_0 h_0 \\ 1^{+-} & 2^{+-} & 0^{+-} \end{array}$$

Mass scale ~ 2 GeV

$$\eta_1 \rightarrow a^+_1 \pi^- \rightarrow (\rho^\circ \pi^+) (\pi^-) \rightarrow \pi^+ \pi^- \pi^+ \pi^-$$

all charged

$$h_0 \rightarrow b^\circ_1 \pi^\circ \rightarrow (\omega \pi^\circ) \gamma\gamma \rightarrow \pi^+ \pi^- \gamma \gamma \gamma \gamma \gamma \gamma$$

many photons

$$h_2 \rightarrow K^+_1 K^- \rightarrow \rho^\circ K^+ K^- \rightarrow \pi^+ \pi^- K^+ K^-$$



strange particles

Final state particles

$\pi^\pm K^\pm \gamma p$

Partial Wave Analysis

- States expected to be broad with multi-particle final states
 - Bump hunting in cross section data not expected to be sufficient
- Need PWA:
 - Identify the J^{PC} of a meson
 - Determine production amplitudes & mechanisms
 - Include polarization of beam, target, spin and parity of resonances and daughters, relative angular momentum



Evidence for Exotic Mesons


<i>State</i>	<i>Processes</i>
$\pi_1(1400) \rightarrow \eta\pi$	$\pi^- N$ Interactions $\bar{p}N$ Annihilations
$\pi_1(1600) \rightarrow \eta'\pi$	
$\pi_1(1600) \rightarrow \rho\pi$	
$\pi_1(1600) \rightarrow b_1\pi$ $\pi_1(1600) \rightarrow f_1\pi$	$\pi^- N$ Interactions
$\pi_1(2000) \rightarrow b_1\pi$ $\pi_1(2000) \rightarrow f_1\pi$	

- Candidates with $J^{PC} = 1^{-+}$
- States are controversial → issues with amplitude analysis
 - Possible leakage due to acceptance or insufficient wave sets
 - Problems with interpretation of line shapes and phases
- Physics interpretation as hybrids instead of qqqq states open to question
- $\pi_1(2000)$ needs confirmation


The GlueX Experiment

- Goal: definitive and detailed mapping of **hybrid meson spectrum**
 - Search for **smoking gun signature** of exotic J^{PC} hybrid mesons
 - Exotics do not mix with $q\bar{q}$ mesons
 - Plans for $s\bar{s}$ and baryon spectroscopy
- Tools for the GlueX Project:
 - Accelerator: 12 GeV electrons, **9 GeV** tagged, linearly polarized **photons** with high flux
 - Detector: **hermiticity**, ability to detect both charged and neutral particles with good resolution
 - **Partial-Wave Analysis**: spin- amplitude of multi-particle final states
 - Computing power: 1 Pb/year data collection, databases, distributed computing, grid services...

12 GeV CEBAF





DOE Generic Project Timeline




Overview of 12 GeV Physics

Hall D exploring origin of confinement by studying exotic mesons




Hall B understanding nucleon structure via generalized parton distributions


Hall C precision determination of valence quark properties in nucleons and nuclei




Hall A short range correlations, form factors, hyper- nuclear physics, future new experiments




Hall D



Architect's rendering of Hall-D Complex



The Hall- D Complex




Coh er ent Bre mss strahlung Beam


- γ -beam energy compromise between polarization and meson mass coverage

- Diamond radiator:

Orientation of crystal planes
→ linearly-polarized γ




**Hadronic
Backgrounds**






The GlueX Detector

- Superconducting Solenoid magnet ~ 2.2 T
- Hermetic detector with large acceptance for charged and neutral particles




Central Drift Chambers

- Track charged particles in central region ($140^\circ < \theta < 20^\circ$)
- 25 radial layers of straw tubes
 - 17 straight layers
 - 4 + 6° stereo layers
 - 4 - 6° stereo layers
 - dE/dx capability for $p < 450$ MeV/c
→ identify protons



Forward Drift Chambers





- Purpose: track forward-going ($\theta < 20^\circ$) charged particles
- Design: 4 packages each containing 6 cathode strip chambers
 - Cathode strip chamber: cathode plane / wire plane / cathode plane
 - Drift chambers with cathode readout
 - Cathode planes divided into strips oriented at $\pm 75^\circ$ with respect to wires
 - Each chamber rotated with respect to its neighbor by 60°
 - Position resolution goal $< 200 \mu\text{m}$




Cathode Strip Chambers

- 3D space point at each wire plane

- Drift time → coordinate away from wire
- Strip centroids → avalanche position along wire



Small-scale prototype




- Preamplifier boards: SIPs
 - Gain $\sim 2.3 \text{ mV}/\mu\text{A}$
 - No pulse shaping
 - No tail-cancellation
- Gas mixture:
 $40\% \text{ Ar} / 60\% \text{ CO}_2$


- Readout for cathode strips: CAEN V792 charge-integrating ADCs
- Readout for sense wires: CAMAC discriminator / F1 TDC

Imaging the wires

- Use centroids on both views to reconstruct wire positions
 - Avalanche occurs near wire → x-positions quantized
 - $x_{\text{wire}} \propto 1/\sqrt{2} (\langle u \rangle + \langle v \rangle)$ using cathode data only
 - Gaussian fits to reconstructed wire positions → resolution






Strip lengths vary between
12.8 cm and 27.8 cm




Barrel Calorimeter

- Photon detection in central region
- Alternating lead + scintillating fiber layers
 - Sampling Fraction = ~ 12%




BCAL in Test Beam




Barrel Calorimeter Configuration

Inner Layers Outer Layers





48 modules (phi sectors)




Barrel Calorimeter




Outer Layers
Inner Layers



Silicon Photomultipliers




- Technology choice for readout of inner layers



Immune to magnetic fields...



SiPM Measurements



$$F = 0.178 \downarrow$$


PDE

$(4.4 \pm 0.5)\%$


$(3.4 \pm 0.5)\%$

- Sensitivity to green wavelengths \Rightarrow good match to BCAL fibers


- Clean photo-electron spectrum observed for 35 μm pixel, $3 \times 3 \text{ mm}^2$ sensor at -20°C
 - Would prefer to run near room temperature...



Time-of-flight Detectors




- Particle identification for forward-going charged tracks
- Two layers of scintillating plastic
 - $250 \times 6 \times 2.54 \text{ cm}^3$ bars
 - ~ 168 channels
 - Timing resolution goal $\sigma < 100 \text{ ps}$




Forward Calorimeter

- Detect photons in the forward direction
- Array of $4 \times 4 \times \text{cm}^2$ lead glass blocks (~ 2800 channels)
 - Crystals already in hand (recycled from E852 and RadPhi)




$$\sigma/E = 7.3\%/\sqrt{E} + 3.6\%$$



DAQ and Trigger

- Trigger rate: 200 kHz, Data rate: 100 MB/ s
- No dead time \Rightarrow pipelined electronics...




Electronics

- Custom electronics in VME- 64X/ VXS



VXS
High Speed
Serial
Backplane







16 channel 250 Msps Flash ADC

Energy Sum Module





Event Simulation

PYTHIA-generated background event



Sample Signal Event

$$\gamma p \rightarrow pb^-_1 \pi^+ \rightarrow p(\omega \pi^-) \pi^+ \rightarrow p\pi^+\pi^-\pi^+\pi^-\gamma\gamma$$



Summary and Outlook

- The 12 GeV upgrade project has passed a major milestone
 - DOE awarded CD- 2 in November (3rd out of 5 CD levels)
- The GlueX experiment is a major part of the upgrade
 - Goal is to map out spectrum of hybrid mesons
 - Major construction project with brand-new hall and detector
 - Detectors are in design and prototype stage
 - ... there's still a lot of work to do!
- CD- 3 (Approval to start construction) is expected next year

We welcome new collaborators!


Hall D Workshop

"Photon- hadron physics with GlueX in Hall D"
Jefferson Lab, March 6- 8, 2008

- Topics include:
 - Chiral anomaly and Primakoff effect
 - Charm production near threshold
 - Exclusive reactions at high momentum transfer
 - Nuclear effects in photo- production
 - Meson and Baryon Spectroscopy
 - Detector upgrades

Additional Slides

Overview of Technical Performance Requirements



Hall D	Hall B	Hall C	Hall A
excellent hermeticity	luminosity 10×10^{34}	energy reach	installation space
polarized	hermeticity	precision	
$E_{\gamma} \sim 8.5 - 9 \text{ GeV}$ photons		11 GeV beamline	
10^8 photons/s		target flexibility	
good momentum/ angle resolution		excellent momentum resolution	
high multiplicity reconstruction		luminosity up to 10^{38}	
		particle ID	

Design Parameters


Capability	Quantity	Range
Charged particles	Coverage	$1^\circ < \theta < 140^\circ$
	Momentum Resolution (5° - 140°)	$\sigma_p/p = 1 - 3\%$
	Position resolution	$\sigma \sim 150\text{-}200 \mu\text{m}$
	dE/dx measurements	$20 < \theta < 140^\circ$
	Time-of-flight measurements	$\sigma_t < 80 \text{ ps}$
	Cerenkov and π/K separation	$\theta < 14^\circ$
	Barrel time resolution	$\sigma_t < (150 + 50/\sqrt{E}) \text{ ps}$
Photon detection	Energy measurements	$2 < \theta < 120^\circ$
	Veto capability	$120 < \theta < 170^\circ$
	LGD energy resolution ($E > 100 \text{ MeV}$)	$\sigma_E/E = (3.6 + 7.3/\sqrt{E})\%$
	Barrel energy resolution ($E > 40 \text{ MeV}$)	$\sigma_E/E = (2 + 5/\sqrt{E})\%$
	LGD position resolution	$\sigma_{x,y} \sim 1 \text{ cm}$
	Barrel position resolution	$\sigma_z \sim 4 \text{ cm}$
DAQ/trigger	Level 1	200 kHz
	Level 3 event rate to tape	15 kHz
	Data rate	100 MB/s
Electronics	Fully pipelined	Flash ADCs, multi-hit TDCs
Photon Flux	Initial: $10^7 \text{ } \gamma/\text{s}$ rate	Final: $10^8 \text{ } \gamma/\text{s}$

Linear Polarization


Linear polarization is:

- ✓ Essential to isolate the production mechanism (M) if X is known
- ✓ A J^P filter if M is known (via a kinematic cut)
- ✓ Degree of polarization is directly related to required statistics
- ✓ Linear polarization separates natural and unnatural parity

States of linear polarization are eigenstates of parity. States of circular polarization are not.



Linear Polarization



For circularly polarized photons: $m = +1$ or $m = -1$ \rightarrow $W(\theta, \phi) \propto |P_{\ell}(\cos \theta)|^2$

For unpolarized photons:
equal mixture of $m = +1$ and $m = -1$ \rightarrow $W(\theta, \phi) \propto |P_{\ell}(\cos \theta)|^2$

For x - linear polarization: \rightarrow $W(\theta, \phi) = |Y_{\ell}^{+1} - Y_{\ell}^{-1}|^2 \propto |P_{\ell}(\cos \theta)|^2 \sin^2 \phi$

For y - linear polarization: \rightarrow $W(\theta, \phi) = |Y_{\ell}^{+1} + Y_{\ell}^{-1}|^2 \propto |P_{\ell}(\cos \theta)|^2 \cos^2 \phi$