

Edmundo S. Barriga, Florida State University

Exotic Mesons





QCD allows for J^{PC} combinations forbidden by the quark model. The goal of GlueX is to unambiguously recognize such states. The table shows states allowed to decay to $\omega\eta$.

l	J^{PC}
0	1^{+-}
1	0^{--} , 1^{--} , 2^{--}
2	1^{+-} , 2^{+-} , 3^{+-}
3	2^{--} , 3^{--} , 4^{--}


Event Selection

We present the whole GlueX Phase I data set. We applied selection cuts to ensure exclusive events.


An important selection was the $2\pi^0$ veto.



We selected ω events. The λ distribution allowed us to approximate the background in the selection. Given the vector nature of ω , events coming from it will linearly rise, while the background will be flat.




$$\lambda = \frac{|\vec{\pi}^+ \times \vec{\pi}^-|^2}{\frac{3}{4} \left(\frac{1}{9} M_{3\pi}^2 - M_\pi^2 \right)^2}$$





Results for $\omega\eta$ Selection

About 485k events survive our selections. These events concentrate in a region known to contain the mesons $\omega(1420)$, $h_1(1595)$, and $\omega(1650)$.



Baryon Contamination

Our sample contains N^* background processes. While some of them, like the $N(1535)$, can be removed with a mass cut, others require future studies.



Summary

- Approximately 485k $\omega\eta$ events from the GlueX Phase I data set
- Work is in process to study N^* background
- Future work will consist of a Partial Wave Analysis

Acknowledgements

GlueX acknowledges the support of several funding agencies and computing facilities: gluex.org/thanks

