

MC Tracking Resolutions


Feb. 22, 2009

David Lawrence JLab

the low-down...


- ~16M single π^+ particles were thrown over a wide range of angles and momenta in order to map out the resolution and acceptance of the GlueX detector.
- The motivation comes from having changed the CDC configuration and having removed the target constraint from the ALT1 fitter
- These results do not include track finding or wire-based fitting. Those tend to increase the left-right ambiguity problem which is not currently handled in the ALT1 fitter.

Track Fitting Efficiency




Relative, total momentum Resolution

Feb. 2008




Feb. 2009




Note: For technical reasons, the plot on the left uses “CONT” option to draw contours while the plot on the right colorizes by bin content.

Relative, total momentum Resolution

Feb. 2008




Feb. 2009




Azimuthal Angular resolution

Feb. 2008




Feb. 2009




Polar Angular Resolution

Feb. 2008




Feb. 2009




π^+ , proton efficiencies

π^+




proton




π^+ , proton Momentum Resolutions

π^+




proton




π^+ , proton ϕ Resolutions

π^+




proton




*White areas are failed fits.
These will be re-fit later with
better binning*

π^+ , proton θ Resolutions

π^+



proton




*White areas are failed fits.
These will be re-fit later with
better binning*

Reconstructed γ Energy Resolution


3.2M photons simulated
and reconstructed

hdgeant



100M photons parametrically
simulated using *hdgeant*
derived resolutions

hdparsim



γ Reconstruction Efficiency

Photon considered found if
reconstructed energy was
within +/- 20% of thrown

hdgeant

hdparsim

